

Volunteer Lake Assessment Program Individual Lake Reports **BROAD BAY, OSSIPEE, NH**

MORPHOMETRIC DATA

TROPHIC CLASSIFICATION KNOWN EXOTIC SPECIES

All sampling data meet water quality standards or thresholds for this parameter.

Watershed Area (Ac.):	224,432	Max. Depth (m):	22.3	Flushing Rate (yr ¹)	34.1	Year	Trophic class	Variable Milfoil
Surface Area (Ac.):	464	Mean Depth (m):	8.3	P Retention Coef:	0.04	1987	OLIGOTROPHIC	
Shore Length (m):	10,600	Volume (m ³):	15,573,500	Elevation (ft):	406	2003	OLIGOTROPHIC	

The Waterbody Report Card tables are generated from the DRAFT 2018 305(b) report on the status of N.H. waters, and are based on data collected from 2008-2017. Detailed waterbody assessment and report card information can be found at www.des.nh.gov/organization/divisions/water/wmb/swqa/index.htm

Designated Use	Parameter	Catego	ry	Comment	Comments				
Aquatic Life	Phosphorus (Total)		Good		Sampling data	is better than the water quality standards or thresholds for this parameter.			
рН			Slightly	Bad	Data periodic	ally exceed water quality standards or thresholds for this parameter by a small margin.			
	Oxygen, Dissolved		Very Go	boc	All sampling c	All sampling data meet water quality standards or thresholds for this parameter.			
	Dissolved oxygen satura		Encoura	aging	Limited data in necessary to the second seco	Limited data for this parameter predicts water quality standards or thresholds are being met; however more dat necessary to fully assess the parameter.			
	Chlorophyll-a		Good	OOD Sampling data is better than the water quality standards or thresholds for this parameter.					
Primary Contact Recreation	ntact Recreation Escherichia coli		No Data		No data for this parameter.				
	Chlorophyll-a		Very Good		All sampling data meet water quality standards or thresholds for this parameter.				
BEACH PRIMARY CONTACT ASSESSMENT STATUS									
BROAD BAY - CAMP HUCKINS BEACH Escher		Escheric	hia coli	Cautionary		Limited data for this parameter predicts exceedance of water quality standards or thresholds; however more data are necessary to fully assess the parameter.			
BROAD BAY - CAMP ROBIN HOOD BEACH Es		Escheric	hia coli Very Good		bd	All sampling data meet water quality standards or thresholds for this parameter.			

LEAVITT BAY - CAMP MARIST BEACH WATERSHED LAND USE SUMMARY

Fry, J., Xian, G., Jin, S., Dewitz, J., Homer, C., Yang, L., Barnes, C., Herold, N., and Wickham, J., 2011. Completion of the 2006 National Land Cover Database for the Conterminous United States, PERS, Vol. 77(9):858-864. For larger image contact NHDES.

Very Good

Escherichia coli Very Good

Escherichia coli



Land Cover Category	% Cover	Land Cover Category	% Cover	Land Cover Category	% Cover
Open Water	3.56	Barren Land	0.64	Grassland/Herbaceous	0.36
Developed-Open Space	2.91	Deciduous Forest	23.33	Pasture Hay	0.85
Developed-Low Intensity	0.74	Evergreen Forest	20.37	Cultivated Crops	0.5
Developed-Medium Intensity	0.24	Mixed Forest	38.49	Woody Wetlands	4.63
Developed-High Intensity	0.04	Shrub-Scrub	2.67	Emergent Wetlands	0.6



VOLUNTEER LAKE ASSESSMENT PROGRAM INDIVIDUAL LAKE REPORTS BROAD BAY, OSSIPEE 2019 DATA SUMMARY

RECOMMENDED ACTIONS: Broad Bay phosphorus and chlorophyll levels were representative of oligotrophic or high quality conditions and the improving chlorophyll levels are encouraging. However, water clarity (transparency) continues to experience years of decline potentially due to periods of darker water color. Continue to measure the relationship between water color and clarity as the increased frequency and intensity of storm events flushes waters rich in dissolved organic matter that imparts a tea color to the water. Conductivity levels, while low, have significantly increased (worsened) since monitoring began. This likely reflects road salt impacts from Rt.25 and residential development in sub-watersheds. Educating watershed residents on proper application of sodium chloride products in winter months and managing stormwater runoff can help reduce the pollutant load. UNH Technology Transfer Center's Road Salt Reduction and NHDES's Watershed Assistance websites are an excellent resource for information on road salt reduction and stormwater management. Keep up the great work!

OBSERVATIONS (Refer to Table 1 and Historical Deep Spot Data Graphics)

- CHLOROPHYLL-A: Chlorophyll levels fluctuated within a low range and were highest in July. Average chlorophyll level decreased slightly from 2018 and was much less than the state median and the threshold for oligotrophic lakes. Historical trend analysis indicates significantly decreasing (improving) chlorophyll levels since monitoring began.
- CONDUCTIVITY/CHLORIDE: Epilimnetic (upper water layer), Metalimnetic (middle water layer) and Hypolimnetic (lower water layer) conductivity levels remained within an average range for NH lakes and approximately equal to the state median. Epilimnetic chloride levels were slightly greater than the state median, yet much less than the state chronic chloride standard. However, historical trend analysis indicates significantly increasing (worsening) epilimnetic conductivity levels since monitoring began.
- COLOR: Apparent color measured in the epilimnion indicated the water fluctuated within a moderately tea colored range and was darkest in May and July.
- TOTAL PHOSPHORUS: Epilimnetic, Metalimnetic and Hypolimnetic phosphorus levels fluctuated within a low range. Average epilimnetic phosphorus level increased from 2018 but remained much less than the state median, and slightly less than the threshold for oligotrophic lakes. Historical trend analysis indicates relatively stable epilimnetic phosphorus levels since monitoring began.
- TRANSPARENCY: Transparency measured with (VS) and without (NVS) the viewscope was below average from May through August and was lowest (worst) in May and July. Transparency then increased (improved) to within a good range in September. Average NVS transparency decreased from 2018 and was approximately equal to the state median. Historical trend analysis indicates significantly decreasing (worsening) transparency is more monitoring bear.
- median. Historical trend analysis indicates significantly decreasing (worsening) transparency since monitoring began.
 TURBIDITY: Epilimnetic and Metalimnetic turbidity levels fluctuated within a low range. Hypolimnetic turbidity level was slightly elevated in September.
- PH: Epilimetic pH levels fluctuated around the low end of the desirable range 6.5-8.0 units. Historical trend analysis indicates stable epilimetic pH levels since montioring began. Metalimetic and Hypolimetic pH levels were slightly less than desirable.

Station Name	Ta	Table 1. 2019 Average Water Quality Data for BROAD BAY - FREEDOM								
	Alk.	Chlor-a	Chloride	Color	Cond.	Total P	Trans.		Turb.	рН
	mg/l	ug/l	mg/l	pcu	us/cm	mg/l	m		ntu	
							NVS	VS		
Epilimnion	5.4	1.61	9	52	49.4	6	3.32	3.76	0.41	6.49
Metalimnion					46.9	6			0.39	6.24
Hypolimnion					45.4	6			0.99	6.05





NH Water Quality Standards: Numeric criteria for specific parameters. Results exceeding criteria are considered a water quality violation. Chloride: > 230 mg/L (chronic) E. coli: > 88 cts/100 mL – public beach E. coli: > 406 cts/100 mL – surface waters Turbidity: > 10 NTU above natural level pH: between 6.5-8.0 (unless naturally occurring)

ters generated from historic lake monitoring data.	
Alkalinity: 4.5 mg/L	
Chlorophyll-a: 4.39 ug/L	
Conductivity: 42.3 uS/cm	
Chloride: 5 mg/L	
Total Phosphorus: 11 ug/L	
Transparency: 3.3 m	
pH: 6.6	

NH Median Values: Median values for specific parame-





This report was generated by the NHDES Volunteer Lake Assessment Program (VLAP). For more information contact VLAP at (603) 271-2658 or sara.steiner@des.nh.gov

HISTORICAL WATER QUALITY TREND ANALYSIS