

Compilation of the 1:100,000-Scale Surficial Geologic Map and Database, Gloucester 100K Quadrangle, New Hampshire

By Michael W. Howley, Joshua A. Keeley, and Jean A. Schwab

New Hampshire Geological Survey
Department of Environmental Services
29 Hazen Drive, Concord, NH 03301

Abstract

This report describes the surficial geology and Quaternary history of the New Hampshire portion of the Gloucester 30x60-minute quadrangle, located in southeastern New Hampshire. Surficial earth materials include unconsolidated sediments (till, sand, gravel, etc.) of glacial and nonglacial origin. Most of these deposits formed during and after the most recent glaciation by the late-Wisconsin Laurentide ice sheet, the last continental glacier to advance over and retreat from the area. Surficial sediments cover the bedrock over hillsides and valley bottoms in the quadrangle and are subject to various land-use considerations. These include sand and gravel extraction, development and protection of ground-water supplies, siting of waste disposal facilities, selection of building sites, and agriculture.

The map compilation for this study was carried out by the authors in support of the STATEMAP cooperative between the New Hampshire Geological Survey and the U. S. Geological Survey (USGS). The geologic map accompanying this report shows the distribution of sedimentary deposits, and indicates their age, composition, and known or inferred origin. It also includes information on the geologic history of the quadrangle, such as features indicating the flow direction of glacial ice. This map provides the basis for the discussion of glacial and postglacial history presented here.

Introduction

This 1:100,000-scale 30x60' surficial geologic map is a compilation of 4 previously mapped 1:24,000-scale 7.5' surficial geologic maps, of which 3 were published by the New Hampshire Geological Survey and 1 was published by the United States Geological Survey (USGS) (Figure 1). The original mapping was completed between 1989 and 2001 by a variety of geologists (Table 1). This compilation project was recommended and approved by the New Hampshire Geologic Resource Advisory Committee to satisfy stakeholders' needs for seamless geologic map data. The map is provided as a high-resolution PDF as well as in ESRI geodatabase format following USGS's data standard, [Geologic Map Schema \(GeMS\)](#). The deliverables include open-source-readable files such as GIS shape files, Microsoft Excel spreadsheets, and text files for all users. Maps, map pamphlets, and GIS data may be requested from the New Hampshire Geological Survey via email to geology@des.nh.gov.

Methods

Previously published and digitized 1:24,000-scale 7.5' map unit GIS polygons were merged and consolidated into 15 surficial units using depositional environment and age categories defined by the

original mapper in an effort to remain consistent with the original data and field relationships. In order to achieve an internally consistent map without redundancies and ambiguities, several slight modifications were made. For example, bedrock polygons originally mapped as “br” were too small and few in number to map at this scale, so they were consolidated into the thin till unit (Ptt), which consists of both thin till and abundant bedrock outcrops.

The original 7.5-minute map publications were in various levels of cartographic completion and not all followed the same standard format. Preparing new publication-ready PDFs for each original 7.5’ map was outside of the scope of the project. However, because the authors realize that map users may compare this compilation to the original publications, the GIS data include a crosswalk between original map units and their new designations in this compilation. That information is stored in the field “Original_MapUnit” in the feature class or shapefile “MapUnitPolys”. The depositional environment information is on the map and stored in the field “DEP_ENV” in the “DescriptionOfMapUnits” table.

The map unit symbols/labels follow a consistent naming scheme. Geologic map unit symbols begin with a capitol letter for age distinction (H, P, or Q for Holocene, Pleistocene, and Quaternary, respectively). Lower case letters stand for depositional environment. Depositional environment categories include alluvial, pluvial (wetland), eolian, glaciolacustrine and glaciofluvial. Subcategories of alluvium include active alluvium (Ha) and stream terraces (Qst). Subcategories of glaciomarine deposits include near-shore (ms) and deltaic (md) deposits. Ice-contact deposits and other fluvial sediments are not differentiated and combined into a glaciofluvial (f) unit. Subcategories of the marine Presumpscot Formation (p) are the sandy facies (ps) deposited in a shallow water environment, and the clay facies (pc) deposited in deeper water.

Many of these maps were published prior to the availability of statewide 1-meter or sub-meter resolution LiDAR data and were mapped on 1:24,000-scale topographic base maps. The statewide Water Well Inventory, maintained by New Hampshire Department of Environmental Services, was used to update the thin till layer systematically across the map using reported depth of bedrock. Steps were taken during the compilation process to assess the accuracy of these geologic map units in comparison to the LiDAR. However, due to the scope of this project, limited steps were taken to align the GIS to the LiDAR. The accuracy of the map unit polygons is generally less than 100 feet, although higher inaccuracies may occur.

Table 1. Summary of compiled 1:24,000-scale geologic surficial maps

Gloucester 30x60-minute sheet			
Quad ID	7.5-minute Quadrangle	Year Published	Author
185	Exeter	2000	Goldsmith/Nelson
186	Hampton	1989	Koteff/Gephart/Shaefer
201	Newburyport West	1991	Goldsmith
202	Newburyport East	1989	Koteff/Gephart/Shaefer




Figure 1 - Index map of compiled 1:24,000-scale geologic surficial maps

Surficial Geologic History of the Gloucester 30x60' Quadrangle, New Hampshire

Surficial deposits include unconsolidated to weakly consolidated deposits overlying bedrock. In the New Hampshire portion of the Gloucester 30'x60' quadrangle, bedrock includes igneous, metaigneous and metasedimentary rocks of Ordovician-Cambrian to Devonian age (Escamilla-Casas, 2003; Hussey et al., 2016). The surficial geology of the Gloucester quadrangle in New Hampshire is primarily the result of glacial and post-glacial activities in the area during the Pleistocene and Holocene epochs, respectively. When the Laurentide Ice Sheet advanced over the area, it mobilized and re-deposited a great quantity of unconsolidated material. Those materials, known as surficial deposits, can be classified into two major categories based on whether the deposits were transported in or deposited by moving water.

Those deposits deposited directly beneath the ice mass or dropped out of suspension into disorganized piles of debris are referred to as glacial till (Pt or Ptt). Those deposits transported by or deposited in water are referred to as stratified drift. Till, a poorly sorted and unstratified mix of clay to boulder size clasts within a relatively fine matrix, is present as a mantle of varying thickness directly above bedrock and underlying other surficial deposits across the map area.

Recent work utilizing annual layers in glacial lake sediments known as varves in the Merrimack River valley, in concert with radiocarbon dating, suggests the Laurentide Ice Sheet began its retreat from southern New Hampshire approximately 16,000 calendar years ago (Ridge et al., 2012). The movement of the Laurentide Ice Sheet across the landscape also shaped the morphology of the area. Streamlined hills of till or drumlins, with long axes aligned with the dominant direction of ice flow, reflect the shaping of the landscape by the moving ice mass. Some exposures of bedrock also retain striations that formed as the ice and entrained materials scraped across the surface. Many such drumlins and striations are shown on the map and record ice movement from the northwest to the southeast.

Moraines, or ridges of till oriented parallel to the front of the ice sheet that indicate ice-margin positions, are present in the Gloucester 30'x60' quadrangle as regularly spaced, low-relief ridges interpreted to be De Geer moraines. The De Geer moraines are interpreted to have formed as products of annual oscillations during recession of the Laurentide Ice Sheet in a glaciomarine setting (Sinclair et al., 2018). The presence of De Geer moraines in southeastern New Hampshire was not recognized until the publication of high-resolution LiDAR data for the Seacoast region of New Hampshire in 2008, shown in Figure 2.

Stratified drift deposits form the major sand and gravel deposits within the Gloucester 30'x60' quadrangle. These deposits were carried away from the receding glacier by meltwater and deposited in marine (ocean) or fluvial (rivers) environments. All stratified drift deposits having undergone some degree of sorting and have become well-rounded by the action of moving water. Coarser materials are generally found near the ice margin (proximal environment) where meltwater energy is the greatest. The further from the ice margin one proceeds, the finer the material that can be carried, so deposits become finer away from the ice margin (distal environment).

Relative sea level fluctuations in the map area during and after the last glaciation were controlled by the interplay of eustatic sea level rise from addition of water from the melting Laurentide Ice Sheet and isostatic rebound of the land surface after the retreat of the Laurentide Ice Sheet. During the Last Glacial Maximum (26,500 to 19,500 years ago). All of the map area was covered by the Laurentide Ice Sheet




Figure 2 – Location map of the Seacoast region of New Hampshire indicating the distribution of De Geer moraines, modified from Figure 1 of Sinclair et al., 2018. Top left inset panel shows the maximum extent of the Laurentide Ice Sheet in North America. Blue shading indicates the marine limit during the late-Pleistocene. Red polygons indicate identified swarms of De Geer moraines in the Seacoast region of New Hampshire. Detail inset (pink) shows high-resolution LiDAR multidirectional hillshade of the regularly spaced, low-relief De Geer moraine swarm in Hampton.

(Clark et al., 2009). As the Laurentide Ice Sheet retreated inland, the ice sheet stayed in contact with the sea. Coastal lowlands in New England were inundated to a present-day elevation of 300 to 320 feet (91 to 98 meters) in the Kittery 100K map area to the north (Tinkham, 2005). Isostatic rebound of the recently deglaciated areas as the ice sheet retreated inland led to a sea-level regression to a low stand of -180 feet (-55 meters) compared to modern sea level (Barnhardt et al., 1995). As relative sea level continued to rise due to the continued addition of meltwater, the marine transgression resumed to present-day sea levels.

In the Gloucester 30'x60' Quadrangle, many stratified deposits were laid down in a marine setting as wave-modified nearshore or beach deposits (Pms) and as deltaic deposits (Pmd) where glacial meltwater entered the synglacial sea and deposited sediments. As the ice receded from the area, the ocean likely maintained contact with the ice across the map area. The late-Pleistocene marine limit in the Gloucester 30'x60' quadrangle presently lies at an elevation of approximately 230 feet (70 meters) above present-day sea level (Ward and Adams, 2001). Only the higher elevation streamlined hills in the

southwest part of the Gloucester 30'x60' quadrangle were above the marine limit. Submarine sand, silt, and clay of the Presumpscot Formation were deposited in the synglacial sea in an estuarine environment as shallow-water deposits (Pp or Pps), or as deep-water deposits (Pp or Ppc).

Non-marine glaciofluvial deposits and eskers (Pf and Pge) are preserved in some areas. Glaciofluvial deposits were deposited directly by glacial streams and rivers above the marine limit. The eskers formed within tunnels in the melting ice mass and remain as steep-sided, narrow piles of coarse-grained debris that are preserved in some areas both above and below the marine limit.

Extensive re-working of the glacial deposits by post-glacial streams and rivers cut stream terraces (Qst) in many locations, which now stand above the level of the existing rivers and streams. Alluvial deposits (Ha) are still being created by the on-going work of present-day streams as they continually erode and deposit unconsolidated material within their floodplains in the quadrangle. Many of the wetlands that exist today (Hw) formed in areas that were previously stream channels or locations where stagnant ice blocks melted away leaving lakes and ponds that continue to be filled in to this day. Deposits near the modern coastline formed by ongoing near-shore processes include sandy beach deposits (Hb) and marine estuary deposits (Hme).

References

- Barnhardt, W.A., Gehrels, W.R., Kelley, J.T., 1995, Late Quaternary relative sea-level change in the western Gulf of Maine: evidence for a migrating glacial forebulge: *Geology* 23: 317-320.
- Clark PU, Dyke AS, Shakun JD et al., 2009, The last glacial maximum: *Science* 325: 710-714.
- Escamilla-Casas, J.C., 2003, [Bedrock geology of the Exeter and Hampton 7.5-minute quadrangles, southeastern New Hampshire](#): New Hampshire Geological Survey, Map Geo-185-024000-BMOF, scale 1:48,000.
- Franzi, David, Wittkop, Chad, and Koteff, Carl, 2005, [Surficial geologic map of the Newmarket quadrangle, Rockingham and Strafford Counties, New Hampshire](#): New Hampshire Geological Survey, Open-File Reports GEO-169-024000-SMAP, scale 1:24,000.
- Goldsmith, Richard, 1991, [Surficial geologic map of the Rockingham County, New Hampshire parts of the Haverhill and Newburyport West quadrangles, New Hampshire and Massachusetts](#): New Hampshire Department of Environmental Services, Open-File Reports NH-91-3, scale 1:24,000.
- Goldsmith, Richard, 2001, [Surficial geologic map of the Exeter quadrangle, Rockingham, County, New Hampshire](#): New Hampshire Department of Environmental Services, scale 1:25,000.
- Hussey, A.M., II, Bothner, W.A., and Thompson, P.J., 2016, [Bedrock geology of the Kittery 1:100,000 quadrangle, Maine and New Hampshire](#): Maine Geological Survey, Geologic Map 16-6, scale 1:100,000.
- Koteff, C., and Pessl, F., Jr., 1985, Till stratigraphy in New Hampshire: Correlations with adjacent New England and Quebec, in Borns, H.W. Jr., LaSalle, P., and Thompson, W.B., eds., Late Pleistocene history of northeastern New England and adjacent Quebec: Geological Society of America Special Paper 197, p. 1-12.

- Koteff, Carl, Goldsmith, Richard, and Gephart, G.D., 1989, [Surficial geologic map of the Dover West quadrangle, Strafford County, New Hampshire](#): U.S. Geological Survey, Open-File Report OF-89-166, scale 1:24,000.
- Koteff, Carl, Gephart, G.D., and Schaefer, J.P., 1989, [Surficial geologic map of the Hampton 7.5 minute quadrangle \(east half of the Exeter 7.5 x 15 minute quadrangle\), New Hampshire-Massachusetts](#): U.S. Geological Survey, Open-File Report OF-89-430, scale 1:24,000.
- Koteff, Carl, Gephart, G.D., and Shaefer, J.P., 1989, [Surficial geologic map of the Newburyport East quadrangle, Massachusetts-New Hampshire](#): New Hampshire Department of Environmental Services, scale 1:24,000.
- Koteff, Carl, 1991, [Surficial geologic map of parts of the Rochester and Somersworth quadrangles, Strafford County, New Hampshire](#): U.S. Geological Survey, Miscellaneous Investigations Series Map I-2265, scale 1:24,000.
- Larson, G.J., 1992, [Surficial geologic map of the Portsmouth and Kittery quadrangles, Rockingham County, New Hampshire](#): New Hampshire Department of Environmental Services, Map SG-6, scale 1:24,000.
- Larson, G.J., and Goldsmith, Richard, 1989, [Surficial geologic map of the Dover East quadrangle in New Hampshire](#): U.S. Geological Survey, Open-File Report OF-89-97, scale 1:24,000.
- Ridge, J.C., Balco, G., Bayless, R.L., et al., 2012, The new North American Varve Chronology: a precise record of southeastern Laurentide Ice Sheet deglaciation and climate, 18.2±12.5 kyr bp, and correlations with Greenland ice core records: *American Journal of Science* 312: 685-722.
- Sinclair, S.N., Licciardi, J.M., Campbell, S.W., and Madore, B.M., 2018, Character and origin of De Geer moraines in the Seacoast region of New Hampshire, USA: *Journal of Quaternary Science* 33: 10.1002/jqs.3017.
- Tinkham, D.J., 2005, [Surficial geologic map of the Milton quadrangle, Strafford County, New Hampshire](#): New Hampshire Department of Environmental Services, Open-File Reports NH-OFM-05-02, scale 1:24,000.
- Ward L.G. and Adams, J.R., 2001, [A preliminary assessment of tidal flooding along the New Hampshire coast: Past, present and future](#): New Hampshire Office of Emergency Management and the Office of State Planning Coastal Program: Concord, NH.